

Risk of Atherosclerotic Cardiovascular Diseases in various Ethnicities of Pakistan

Tariq Ashraf¹, Asif Nadeem²,
Shakil Sarwar³, Musa Karim⁴

ABSTRACT

Objective: To estimate the risk of atherosclerotic cardiovascular diseases (ASCVD) in various ethnicities of Pakistan using Astronaut Cardiovascular Health and Risk Modification (Astro-CHARM) risk calculator and the Pooled Cohort Equations (PCE).

Methods: Individuals of both gender between 40 to 65 years of age without pre-existing coronary artery disease from residents of Karachi were recruited through snowball sampling technique. Study was conducted at the National Institute of Cardiovascular Diseases, Karachi, Pakistan during January to June 2019. Ethnicity of the participants was categorized based on their mother tongue. Ten-year risk of ASCVD event was estimated using Astro-CHARM Calculator and PCE.

Results: Mean age of a total of 386 individuals was 49(±7.1) years and 45.3% (175) were females. Urdu-speaking individuals were 33.4%(129) of the sample, while, 18.4%(71) Pushtoons, 14%(54) Sindhi, 13%(50) Punjabi, 2.6%(10) Balochi, and remaining 18.7%(72) were of other ethnicities. High risk (≥7.5%) individuals were 20.7% (80/386) as per PCE and 11.1% (43/386) as per Astro-CHARM. As per Astro-CHARM, Sindhis' had the least risk of ten-years ASCVD event among all the ethnicities, while, Urdu-speakers' had the highest risk with mean rank of 145.18 vs. 216.50, p-value=0.001.

Conclusion: A significant 10-years risk of first ASCVD event was observed in our population. ASCVD risk is alarmingly high in some ethnicities, such as Urdu-speaking, owing to the increased prevalence of traditional modifiable risk factors, such as diabetes and smoking.

KEYWORDS: Atherosclerotic Cardiovascular Diseases, Risk Assessment, Pooled Cohort Equations, Ethnicities, Pakistan, Astro-CHARM, Primary Prevention.

doi: <https://doi.org/10.12669/pjms.36.6.2618>

How to cite this:

Ashraf T, Nadeem A, Sarwar S, Karim M. Risk of Atherosclerotic Cardiovascular Diseases in various Ethnicities of Pakistan. *Pak J Med Sci.* 2020;36(6):1158-1164. doi: <https://doi.org/10.12669/pjms.36.6.2618>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Tariq Ashraf
2. Asif Nadeem
3. Shakil Sarwar
4. Musa Karim
- 1,4: National Institute of Cardiovascular Diseases (NICVD), Karachi, Pakistan.
- 2,3: Combined Military Hospital (CMH) Malir, Karachi, Pakistan.

Correspondence:

Prof. Dr. Tariq Ashraf
National Institute of Cardiovascular Diseases (NICVD),
Karachi, Pakistan.
E-mail: tariqash45@gmail.com

- * Received for Publication: April 3, 2020
- * 1st Revision Received: May 6, 2020
- * 2nd Revision Received: June 26, 2020
- * Final Revision Accepted: * June 28, 2020

INTRODUCTION

About one-fourth of the world population, living in the South Asian region, are at an increased risk of atherosclerotic cardiovascular diseases (ASCVD).^{1,2} Worlds' efforts to curtail the burden of ASCVD is paying off with the decline in the global burden of diseases, however, for the South Asian region it is on its rise.³ The increased risk of coronary artery diseases (CAD) among the South Asian population is ingrained in pathophysiological as well as lifestyle-related factors.⁴ A meta-analysis highlighted that the cross-sectional area of the coronary artery is smaller in South Asian population which increases the risk

of CAD progression and development.⁵ Although, the biological mechanism of development of ASCVD is complex but increased prevalence of established risk factors in this population can be partly attributed to the increased risk of ASCVD.¹

Pakistani population comprises of over 200 million inhabitants of the South Asian region. According to a recent study, more than 30% of the Pakistani individuals were at high risk ($\geq 7.5\%$) of 10 years ASCVD events, smoking, diabetes (type II), hypertension, and high cholesterol were the most prevalent modifiable risk factors.⁶

Various risk assessment tools and clinical calculators have been developed for the assessment of ASCVD risk such as Framingham risk score, Reynolds risk score, systematic coronary risk evaluation, World Health Organization risk charts, Lancet chronic diseases risk charts, European system for cardiac operative risk evaluation (EuroSCORE), etc., but their clinical utility is limited due to poor concordance and predictability.⁷ According to the recent ACC/AHA primary prevention guidelines,⁸ individuals with an estimated risk of $\geq 7.5\%$, calculated based on the Pooled Cohort Equation (PCE), with coronary artery calcium (CAC) measured ≥ 100 Agatston units statin therapy should be initiated. The Astronaut Cardiovascular Health and Risk Modification (Astro-CHARM) risk calculator is an improvement over PCE with the addition of CAC and C-reactive protein (CRP).⁹

Hence, risk assessment is a vital step in implementing and developing preventive strategies. Available data elaborating risk of ASCVD for the Pakistani population and its ethnic breakup are based on traditional risk factors based calculators and data regarding CAC for our population is not available. The purpose of this study was to estimate the risk of atherosclerotic cardiovascular diseases (ASCVD) in various ethnicities of Pakistan using Astro-CHARM risk calculator and the Pooled Cohort Risk Assessment Equations.

METHODS

This study was conducted at the National Institute of Cardiovascular Diseases (NICVD), Karachi, Pakistan during January to June 2019. Approval of the Ethical Review Committee (ERC-29/2018) was obtained and written informed consent was obtained from all the participants regarding the publication of their data while maintaining confidentiality. Both males and females of 40 to

65 years of age without pre-existing CAD were recruited using snowball sampling technique. Geographic coverage and ethnic distribution of the recruited individuals were ensured, detailed research methodology is presented elsewhere.¹⁰

The demographic data (gender, mother tongue, age (years), height (cm), and weight (kg)), lipid profile (total, HDL, and LDL cholesterol), systolic blood pressure measurement, and C-reactive protein (CRP) were obtained. History of the patients regarding traditional risk factors were obtained as per user guide of Astro-CHARM Calculator. Coronary computed tomography angiography (CCTA) was performed in all individuals to obtain coronary artery calcium (CAC) score (Agatston units).

Pakistani populations comprises of diverse racial and ethnical clusters dwelling in urban and rural areas. Ethnic distribution in rural areas are mostly outlined by administrative as well as geographic boundaries, nevertheless, the distribution in urban areas, especially in major cities, is not very well decisive.¹¹ As, mother tongue (native language) is considered to be a strong surrogate of ethnicity, therefore, ethnicity of the participants was categorized as Sindhi, Punjabi, Pushtoon, Balochi, Urdu speaking (Muhajirs), and others (Sirayki, Bangali, Kashmiri, Gilgiti, etc.) based on the mother tongue.¹² Ten-year risk of atherosclerotic cardiovascular diseases (ASCVD) event was estimated using Astro-CHARM Calculator⁹ (available at www.astrocharm.org) and Pooled Cohort Equation (PCE) Risk Assessment Equations¹³ (available at <https://clincalc.com/cardiology/ascvd/pooledcohort.aspx>). Variables included in the calculation of PCE and Astro-CHARM risk calculators are presented in Table-I.

The race variable was entered as "others" for Astro-CHARM Calculator and "white or other" for PCE. Participants were stratified into four groups based CAC score, no risk, low risk, medium risk, and high risk group with CAC score of 0 Agatston Units, 1 to 99 Agatston Units, 100 to 399 Agatston Units, and ≥ 400 Agatston Units respectively. Individuals with $\geq 7.5\%$ of 10-years ASCVD risk based on Astro-CHARM or PCE were stratified as high risk.

All the data were collected on a predefined structural proforma entered on a data entry screen, developed using CsPro 7.0 by the U.S. Census Bureau and ICF International. Serpro S.A. Data were analyzed using International Business Machines (IBM) Statistical Package for

Table-I: Variables included in the calculation of risk calculators.

Variables	Calculator	
	Astro-CHARM	Pooled Cohort Equation (PCE)
Gender	Included	Included
Age	Included	Included
Race	Included	Included
Total cholesterol	Included	Included
HDL cholesterol	Included	Included
Systolic blood pressure	Included	Included
Hypertension	Included	Included
Diabetes	Included	Included
Smoking	Included	Included
Family history of heart attack	Included	Not included
Coronary Artery Calcium	Included	Not included
C-reactive protein (CRP)	Included	Not included

the Social Sciences (SPSS), Version 21.0. Mean (\pm standard deviation), median [interquartile range (IQR)], minimum, and maximum of the ten-year atherosclerotic cardiovascular diseases (ASCVD) event risk were calculated for all the ethnicities. The Kolmogorov-Smirnov test (KS test) was applied for the calculated score of Astro-CHARM and PCE and hypothesis of normality of the distribution of calculated score was rejected. The Kruskal-Wallis test followed by pairwise comparisons were made to assess the calculated score by various ethnic groups. The p -value ≤ 0.05 was the set criteria for the statistical significance.

RESULTS

In this study 386 individuals with mean age of 49 (± 7.1) years without pre-existing CAD were included, out of which 45.3% (175) were females, 62.4% (241) were 40 to 50 years of age. The traditional risk factors of the cardiovascular diseases were as following; 45.6% (176) individuals were on antihypertensive treatment, 15.8% (61) were taking hyperglycemic treatments, 14.2% (55) were active smokers, and 30.6% (118) of these individuals had history of heart attack in their first degree relatives. The ethnic breakup was as follows; 33.4% (129) sample consisted of Urdu speaking individuals, 18.4% (71) were Pushtoon, 14% (54) were Sindhi, 13% (50) were Punjabi, 2.6% (10) were Balochi, and remaining 18.7% (72) of the individuals were of other ethnicities comprises of Sirayki, Bangali, Kashmiri, Gilgiti, etc. Demographic characteristics and distribution of traditional risk factors among the study participants by various ethnicities are presented in Table-II.

The summary of ten-year risk of atherosclerotic cardiovascular diseases (ASCVD) event estimated using Astro-CHARM and Pooled Cohort Equation (PCE) for the ethnicities are presented in Table-III. The Kruskal-Wallis test revealed that the distribution of risk calculated using the Pooled Cohort Equation (PCE) is same across the ethnicities with p -value of 0.061 (Chi-square = 10.538, degrees of freedom = 5). While, the distribution of risk calculated using Astro-CHARM calculator is not same across the ethnicities with p -value of 0.003 (Chi-square = 18.301, degrees of freedom = 5). The pairwise

Table-II: Demographic characteristics and distribution of traditional risk factors among the study participants by various ethnicities.

Characteristics	Sindhi	Punjabi	Balochi	Pushto	Urdu	Others	p -value
	N=54	N=50	N=10	N=71	N=129	N=72	
Age (years)	47 (± 7.3)	49.5 (± 7.9)	46.9 (± 8.7)	48.3 (± 6.6)	50.3 (± 6.8)	48.8 (± 6.9)	0.019
40 to 50 years	42 (77.8%)	31 (62%)	7 (70%)	46 (64.8%)	67 (51.9%)	48 (66.7%)	0.029
51 to 65 years	12 (22.2%)	19 (38%)	3 (30%)	25 (35.2%)	62 (48.1%)	24 (33.3%)	
Gender							
Male	33 (61.1%)	28 (56%)	3 (30%)	31 (43.7%)	73 (56.6%)	43 (59.7%)	0.168
Female	21 (38.9%)	22 (44%)	7 (70%)	40 (56.3%)	56 (43.4%)	29 (40.3%)	
Medical History							
Hypertension	26 (48.1%)	23 (46%)	5 (50%)	39 (54.9%)	47 (36.4%)	36 (50%)	0.177
Diabetes	2 (3.7%)	6 (12%)	2 (20%)	14 (19.7%)	26 (20.2%)	11 (15.3%)	0.098
Currently Smoke	6 (11.1%)	7 (14%)	1 (10%)	6 (8.5%)	20 (15.5%)	15 (20.8%)	0.385
FHxHA	10 (18.5%)	19 (38%)	4 (40%)	21 (29.6%)	49 (38%)	15 (20.8%)	0.035

FHxHA = family history of heart attack.




Fig.1: The pairwise comparisons of ten-year risk of atherosclerotic cardiovascular diseases (ASCVD) by ethnicities.

comparison of the calculated risk among the various ethnicities showed that the "Sindhi" had the least risk of ten years ASCVD event among all the ethnicities, while, "Urdu Speaking" had the highest risk of ten years ASCVD event with mean rank of 145.18 vs. 216.50, p-value = 0.001. The pairwise comparisons of ten-year risk of atherosclerotic cardiovascular diseases (ASCVD) by ethnicities are presented in Fig.1.

According to PCE, 20.7% (80/386) of individuals were at high risk ($\geq 7.5\%$) of 10-years ASCVD event. Burden of high risk of 10-years ASCVD event by the ethnicities are as following; rates of high risk were 11.1% (6/54) for Sindhi, 24% (12/50) for Punjabis, 10% (1/10) for Balochi, 16.9% (12/71) for Pushtoon, 24.8% (32/129) for Urdu speaking individuals and 23.6% (17/72) for other ethnicities. Similarly, according to Astro-CHARM, 11.1% (43/386) were high risk ($\geq 7.5\%$) individuals, rates of high risk were 1.9% (1/54) for Sindhi, 14% (7/50) for Punjabi, 0% (0/10) for Balochi, 7% (5/71) for Pushtoon, 18.6% (24/129) for Urdu speaking individuals and 8.3% (6/72) for other ethnicities.

Table-III: The summary of ten-year risk of atherosclerotic cardiovascular diseases (ASCVD) event estimated using Astro-CHARM and Pooled Cohort Equation (PCE) for the ethnicities.

Ethnicities	Base		10-years ASCVD Event Risk (%)		
	N	Mean (\pm SD)	Median [IQR]	Range	Mean Rank
Astro-CHARM					
Total	386	3.57 (\pm 4.62)	1.9 [4 - 1.1]	40.3 - 0.2	-
Sindhi	54	1.88 (\pm 1.59)	1.4 [2.2 - 0.9]	9 - 0.3	145.18
Punjabi	50	4.08 (\pm 5.43)	1.9 [5.4 - 1.3]	31.4 - 0.5	201.68
Balochi	10	2.13 (\pm 1.56)	2 [2.6 - 1]	5.9 - 0.6	166.55
Pushto	71	2.74 (\pm 2.82)	1.8 [3.5 - 1]	17.7 - 0.2	177.83
Urdu	129	4.62 (\pm 5.78)	2.4 [5.7 - 1.2]	40.3 - 0.5	216.50
Others	72	3.6 (\pm 4.42)	2.45 [4.05 - 1.15]	25 - 0.4	202.05
Pooled Cohort Equation (PCE)					
Total	386	4.96 (\pm 5.79)	2.8 [6.2 - 1.3]	43.8 - 0.2	-
Sindhi	54	2.99 (\pm 2.77)	2.1 [4.2 - 1.1]	12.1 - 0.2	156.89
Punjabi	50	5.16 (\pm 5.84)	2.95 [6.8 - 1.4]	25.1 - 0.5	197.47
Balochi	10	4.03 (\pm 5.06)	2.45 [5.4 - 1]	17.5 - 0.5	168.65
Pushto	71	4.15 (\pm 4.52)	2.7 [5.7 - 1.1]	26.1 - 0.4	182.21
Urdu	129	5.83 (\pm 6.36)	3.7 [7 - 1.7]	33.4 - 0.3	210.43
Others	72	5.67 (\pm 7.1)	3 [7 - 1.6]	43.8 - 0.5	202.45

Table-IV: Distribution of the coronary artery calcium (CAC) score for the various ethnicities.

CAC (Agatston Units)	Sindhi	Punjabi	Balochi	Pushto	Urdu	Others
	N=54	N=50	N=10	N=71	N=129	N=72
No risk - (0)	46 (85.2%)	34 (68%)	8 (80%)	58 (81.7%)	81 (62.8%)	55 (76.4%)
Low risk - (1 to 99)	6 (11.1%)	10 (20%)	2 (20%)	11 (15.5%)	36 (27.9%)	14 (19.4%)
Moderate risk - (100 to 399)	2 (3.7%)	4 (8%)	0 (0%)	1 (1.4%)	9 (7%)	3 (4.2%)
High risk - (≥ 400)	0 (0%)	2 (4%)	0 (0%)	1 (1.4%)	3 (2.3%)	0 (0%)

CAC = coronary artery calcium score.

No coronary artery calcium was observed in 73.1% (282) of the individuals, 20.5% (79) were in the low-risk groups with CAC score of 1 to 99 Agatston Units, 4.9% (19) were in moderate-risk groups with CAC score between 100 and 399 Agatston Units, and remaining 1.6% (6) individual were in high-risk group with CAC score ≥ 400 Agatston Units. Distribution of the coronary artery calcium (CAC) score for the various ethnicities are presented in Table-IV.

DISCUSSION

This is the first study from Pakistan, or even in the South Asian region, reporting 10 years ASCVD risk estimates using Astro-CHARM calculator. Pakistani inhabitants are multicultural with each sub-culture and ethnicity having distinctive lifestyle, socio-economic, and cultural values.¹⁴ Therefore, it was deemed important to assess the risk of atherosclerotic cardiovascular diseases among the various ethnicities dwelling in Pakistan. Low to middle socio-economic class dominates the Pakistani population landscape with a diverse cultural and ethnic breakup. Studies in the past reported great variations in the incidence and risk of ASCVD among the various ethnic breakups of Pakistan.^{10,12,15}

We observed that our Urdu speaking community (commonly known as Muhajirs) is at increased risk of development of ASCVD event in the following 10 years period and presence of coronary artery calcium was also higher in this community. Diabetes, smoking, and family history of heart attack was relatively more common in Urdu speaking communities as compared to the other ethnic groups. In this study individuals of Sindhi ethnic origin were found to have least estimated risk of 10 years ASCVD event with least diabetics of all the other ethnicities. Punjabi are the second most risky ethnicity with relatively high prevalence of traditional risk factors.

These findings are aligned with the findings of a past study by Ashraf T et al.¹², in their study of

non-atherosclerotic Pakistani individuals, 44.87% of Urdu speaking had $\geq 7.5\%$ risk of atherosclerotic event in 10 years followed by Punjabi with 19.23% and high-risk individuals were in the smallest ratio for Sindhi ethnicity. This study along with other studies in this population,^{6,8,10,12} estimates a significant risk of ASCVD in Pakistani population, and for subgroups, such as Urdu speaking and Punjabi, it is alarmingly high.

The coronary artery calcium (CAC) is one of the robust independent predictor and a novel surrogate marker of subclinical ASCVD.¹⁶⁻¹⁸ Progression in the repeated assessment of CAC on computed tomography (CT) scan was reported to be associated with the progression and incidence of CAD and mortality.¹⁹⁻²¹ For the adult population risk of fatal and nonfatal cardiac events observed to be associated with the presence of CAC.^{22,23} Considering the strong predictability, traditional risk factors based risk assessment calculators¹³ were upgraded by integrating CAC score with the conventional risk factors to form a robust risk stratification modality for the ASCVD,⁹ and the recent 2019 ACC/AHA guidelines for the primary prevention of CAD recommended ASCVD risk calculation and CAC measurement to guide the statin therapy.⁸

A systematic risk assessment and initiations of preventive measures including lifestyle modification and lipid-lowering medications are the need of the hour for our population. However, neither Astro-CHARM calculator nor PCE are validated for this population, therefore, future research efforts are also needed to optimize the risk assessment and stratification methods.

A popular saying of "diversity is strength" doesn't seem to favor the Pakistani population when it comes to ASCVD risk assessment and prevention. With a bigger aim of reduction of ASCVD burden at national as well as regional level, it is important to study the sub-clinical and clinical factors in the various ethnicities at its native

grounds and propagate the targeted preventive strategies. Secondly, awareness campaigns and assurance of adaptation of evidence-based medicine and guidelines need to be ensured not only in cardiology clinics but also at the general practitioners' office in urban as well as far-flung rural areas of Pakistan.

Limitations of the study: The Snowball sampling technique adopted for participant recruitment is the limitation of the study, secondly, ethnic distribution of study sample was not as per population distribution of Pakistan, hence, the generalizability of results is limited. Categorization of ethnicities based on mother tongue may does not reflect true ethnicity especially for "Urdu speaking".

CONCLUSION

In the present study, we observed a significant estimated 10 years risk of first atherosclerotic cardiovascular disease (ASCVD) in Pakistani individuals of diverse ethnic backgrounds without preexisting cardiovascular diseases. The risk of ASCVD is alarmingly high for some of the ethnic groups, such as Urdu speaking, owing to the increased prevalence of traditional modifiable risk factors, such as diabetes and smoking.

Acknowledgement: Authors are thankful to the Medical Affairs division of Getz Pharma for providing financial support for CT angiography and lab assessments of the study participants. The authors wish to acknowledge the support of the staff member of the clinical research department of the National Institute of Cardiovascular Diseases (NICVD), Karachi Pakistan.

Disclaimer: None to declare.

Conflict of interest: None to declare.

Funding disclosure: Computed Tomography (CT) angiography and laboratory assessments were financed by Medical Affairs division of Getz Pharma.

REFERENCES

1. Volgman AS, Palaniappan LS, Aggarwal NT, Gupta M, Khandelwal A, Krishnan AV, et al. Atherosclerotic cardiovascular disease in South Asians in the United States: epidemiology, risk factors, and treatments: a scientific statement from the American Heart Association. *Circulation*. 2018;138(1):e1-34. doi: 10.1161/CIR.0000000000000580
2. George J, Mathur R, Shah AD, Pujades-Rodriguez M, Denaxas S, Smeeth L, et al. Ethnicity and the first diagnosis of a wide range of cardiovascular diseases: Associations in a linked electronic health record cohort of 1 million patients. *PLoS One*. 2017;12(6):e0178945. doi: 10.1371/journal.pone.0178945
3. Ahmed ST, Rehman H, Akeroyd JM, Alam M, Shah T, Kalra A, et al. Premature coronary heart disease in South Asians: burden and determinants. *Curr Atheroscler Rep*. 2018;20(1):6. doi: 10.1007/s11883-018-0706-1
4. Nair M, Prabhakaran D. Why do South Asians have high risk for CAD? *Glob Heart*. 2012;7(4):307-314. doi: 10.1016/j.heart.2012.09.001
5. Basu D, Pokharel H, Tiwari A. South Asian ethnicity associated with narrow coronary arteries: a meta-analysis. *J Am Coll Cardiol*. 2018;71(11-Suppl):A203. doi: 10.1016/S0735-1097(18)30744-7
6. Hassan K, Mohyidin B, Fawwad A, Waris N, Iqbal S, Jawaid M. Predicting the risk of atherosclerotic cardiovascular disease (ASCVD) in Pakistani population. *Clin Epidemiol Glob Health*. 2019;7(2):184-187. doi: 10.1016/j.cegh.2018.04.002
7. Bazo-Alvarez JC, Quispe R, Peralta F, Poterico JA, Valle GA, Burroughs M, et al. Agreement between cardiovascular disease risk scores in resource-limited settings: evidence from 5 Peruvian sites. *Crit Pathw Cardiol*. 2015;14(2):74-80. doi: 10.1097/HPC.0000000000000045
8. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: executive summary: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2019;74(10):1376-1414. doi: 10.1016/j.jacc.2019.03.009
9. Khera A, Budoff MJ, O'Donnell CJ, Ayers CA, Locke J, de Lemos JA, et al. Astronaut Cardiovascular Health and Risk Modification (Astro-CHARM) coronary calcium atherosclerotic cardiovascular disease risk calculator. *Circulation*. 2018;138(17):1819-1827. doi: 10.1161/CIRCULATIONAHA.118.033505
10. Ashraf T, Mengal MN, Muhammad AS, Tareen AK, Khan MN, Kazmi KA, et al. Ten years risk assessment of atherosclerotic cardiovascular disease using Astro-CHARM and pooled cohort equation in a south Asian sub-population. *BMC Public Health*. 2020;20(1):403. doi: 10.1186/s12889-020-08472-4
11. Mushtaq M. Managing Ethnic diversity and federalism in Pakistan. *Eur J Sci Res*. 2009;33(2):279-94.
12. Ashraf T, Aamir KF, Achakzai AS, Mengal N, Karim MT. Burden of atherosclerosis risk among different ethnicity: a hidden health challenge. *J Ayub Med Coll Abbottabad*. 2016;28(4):788-792.
13. Lloyd-Jones DM, Leip EP, Larson MG, d'Agostino RB, Beiser A, Wilson PW, et al. Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation*. 2006;113(6):791-798. doi: 10.1161/CIRCULATIONAHA.105.548206
14. Shah SA, Amjad S. Cultural diversity in Pakistan: national vs provincial. *Mediterr J Soc Sci*. 2011;2(2):331-344.
15. Ashraf T, Achakzai AS, Farooq F, Memon MA, Mengal N, Abbas KY, et al. Estimating risk of atherosclerotic cardiovascular diseases in non-atherosclerotic Pakistani patients: study conducted at National Institute of Cardiovascular Diseases, Karachi, Pakistan. *J Pak Med Assoc*. 2017;67(4):494-498.

16. Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med*. 2008;358(13):1336-1345. doi: 10.1056/NEJMoa072100
17. Polonsky TS, McClelland RL, Jorgensen NW, Bild DE, Burke GL, Guerci AD, et al. Coronary artery calcium score and risk classification for coronary heart disease prediction. *JAMA*. 2010;303(16):1610-1616. doi: 10.1001/jama.2010.461
18. Yeboah J, McClelland RL, Polonsky TS, Burke GL, Sibley CT, O'Leary D, et al. Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA*. 2012;308(8):788-795. doi: 10.1001/jama.2012.9624
19. Shemesh J, Apter S, Stoler D, Itzchak Y, Motro M. Annual progression of coronary artery calcium by spiral computed tomography in hypertensive patients without myocardial ischemia but with prominent atherosclerotic risk factors, in patients with previous angina pectoris or healed acute myocardial infarction, and in patients with coronary events during follow-up. *Am J Cardiol*. 2001;87(12):1395-1397. doi: 10.1016/s0002-9149(01)01561-2
20. Budoff MJ, Hokanson JE, Nasir K, Shaw LJ, Kinney GL, Chow D, et al. Progression of coronary artery calcium predicts all-cause mortality. *JACC Cardiovasc Imaging*. 2010;3(12):1229-1236. doi: 10.1016/j.jcmg.2010.08.018
21. Budoff MJ, Young R, Lopez VA, Kronmal RA, Nasir K, Blumenthal RS, et al. Progression of coronary calcium and incident coronary heart disease events: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol*. 2013;61(12):1231-1239. doi: 10.1016/j.jacc.2012.12.035
22. Carr JJ, Jacobs DR, Terry JG, Shay CM, Sidney S, Liu K, et al. Association of coronary artery calcium in adults aged 32 to 46 years with incident coronary heart disease and death. *JAMA Cardiol*. 2017;2(4):391-399. doi: 10.1001/jamacardio.2016.5493
23. Thomas IC, Shiao B, Denenberg JO, McClelland RL, Greenland P, De Boer IH, et al. Association of cardiovascular disease risk factors with coronary artery calcium volume versus density. *Heart*. 2018;104(2):135-143. doi: 10.1136/heartjnl-2017-311536

Authors' Contribution:

TA: Conceived the idea, **TA, MK & AN** Designed the study, **TA, AN & SS:** Collected the data, **TA & MK:** Drafted the manuscript, **AN & SS:** Critically reviewed the manuscript.
TA: Agrees to be accountable for all aspects of work ensuring integrity and accuracy.

Retraction Announcement

The following manuscript has been retracted from November-December 2017 issue on the request of the authors who stated that "after publication, their group found that it was difficult to repeat the results. We believe that there may be some flaws or operational loopholes, hence we would like to retract this paper." - *Editor*

Retraction in: Pak J Med Sci 2017;33(6):1468-1472. **DOI:** <https://doi.org/10.12669/pjms.336.13488>

Link: <http://pjms.com.pk/index.php/pjms/article/view/13488>

Therapeutic effects of Pulpotomy and Pulpectomy on deciduous molars with deep caries

Yuxiang Tang¹, Wantian Xu²

Retracted on July 7, 2020